

## APPLICATION GUIDE:

### CALAMANSI

(*Citrofortunella microcarpa*)



#### Dilution:

- 1:100
- Mix 1 liter of SG7 with 100 liters of water.

#### Dosage:

- 1 ha = 2 liters of SG7 + 186 liters of water

#### Total SG7:

- 18–20 liters per hectare

#### IMPORTANT!

- Calamansi seedlings are ready for transplanting when they have grown to around 3 feet tall (approximately 1–12 months old).
- Transplant seedlings around June or July to ensure that water for irrigation is in adequate supply.
- Apply SG7 early in the morning, between 5–9 a.m. (maximum stomatal opening), or late in the afternoon, between 4–6 p.m. (extended stomatal opening).
- Use unchlorinated water when mixing SG7. (Chlorine can kill microbes in the soil that may be beneficial to plants.)
- As soon as fruits start to develop from the flowers, stop application of SG7 to ensure that fruits are free of residual deposits.
- If cropping started from seedlings, the plant will reach its flowering stage in 18–24 months and its fruiting stage in another 10–12 months.

#### Method & Frequency of Application:

##### 1<sup>st</sup> APPLICATION:

1–3 days before transplanting

- Dosage: 2 liters per hectare or 160 ml per 16 liters tank load
- Method: Apply directly to the soil (moist to dry)

##### 2<sup>nd</sup> APPLICATION:

10–15 days after transplanting

- Dosage: 2 liters per hectare or 160 ml per 16 liters tank load
- Method: Apply directly to the plant

##### SUCCEEDING APPLICATION:

30 days after previous application with a maximum of 6–7 applications

- Dosage: 2 liters per hectare or 160 ml per 16 liters tank load
- Method: Apply directly to the plant

##### LAST APPLICATION:

10–15 days before flowering (approximately 9–10 months after transplanting)

- Dosage: 2 liters per hectare or 160 ml per 16 liters tank load
- Method: Apply directly to the plant